The background of the slide features a complex, abstract molecular structure. It consists of numerous small, glowing spheres in shades of red, white, and blue, interconnected by thin, translucent lines. The overall effect is a sense of dynamic, interconnectedness, typical of scientific or technological branding.

MEDIVIR PRESENTING AT REDEYE LIFE SCIENCE DAY STOCKHOLM, NOVEMBER 21, 2018

MEDIVIR
SCIENCE WORKING WONDERS

Important notice

You must read the following before continuing. The following applies to this document and the information provided in this presentation by Medivir AB (publ) (the "Company") or any person on behalf of the Company and any other material distributed or statements made in connection with such presentation (the "Information"), and you are therefore advised to carefully read the statements below before reading, accessing or making any other use of the Information. In accessing the Information, you agree to be bound by the following terms and conditions.

The Information does not constitute or form part of, and should not be construed as, an offer of invitation to subscribe for, underwrite or otherwise acquire, any securities of the Company or a successor entity or any existing or future subsidiary or affiliate of the Company, nor should it or any part of it form the basis of, or be relied on in connection with, any contract to purchase or subscribe for any securities of the Company or any of such subsidiaries or affiliates nor shall it or any part of it form the basis of or be relied on in connection with any contract or commitment whatsoever. Specifically, this presentation does not constitute a "prospectus" within the meaning of the U.S. Securities Act of 1933, as amended.

The Information may not be reproduced, redistributed, published or passed on to any other person, directly or indirectly, in whole or in part, for any purpose. The Information is not directed to, or intended for distribution to or use by, any person or entity that is a citizen or resident of, or located in, any locality, state, country or other jurisdiction where such distribution or use would be contrary to law or regulation or which would require any registration or licensing within such jurisdiction. The Information is not for publication, release or distribution in the United States, Australia, Canada or Japan, or any other jurisdiction in which the distribution or release would be unlawful.

All of the Information herein has been prepared by the Company solely for use in this presentation. The Information contained in this presentation has not been independently verified. No representation, warranty or undertaking, express or implied, is made as to, and no reliance should be placed on, the fairness, accuracy, completeness or correctness of the Information or the opinions contained herein. The Information contained in this presentation should be considered in the context of the circumstances prevailing at that time and will not be updated to reflect material developments which may occur after the date of the presentation. The Company may alter, modify or otherwise change in any manner the content of this presentation, without obligation to notify any person of such revision or changes.

This presentation may contain certain forward-looking statements and forecasts which relate to events and depend on circumstances that will occur in the future and which, by their nature, will have an impact on the Company's operations, financial position and earnings. The terms "anticipates", "assumes", "believes", "can", "could", "estimates", "expects", "forecasts", "intends", "may", "might", "plans", "should", "projects", "will", "would" or, in each case, their negative, or other variations or comparable terminology are used to identify forward-looking statements. There are a number of factors that could cause actual results and developments to differ materially from those expressed or implied in a forward-looking statement or affect the extent to which a particular projection is realized. Factors that could cause these differences include, but are not limited to, implementation of the Company's strategy and its ability to further grow, risks associated with the development and/or approval of the Company's products candidates, ongoing clinical trials and expected trial results, the ability to commercialize existing and any future products, technology changes and new products in the Company's potential market and industry, the ability to develop new products, the impact of competition, changes in general economy and industry conditions and legislative, regulatory and political factors. While the Company always intends to express its best judgment when making statements about what it believes will occur in the future, and although the Company bases these statements on assumptions that it believes to be reasonable when made, these forward-looking statements are not a guarantee of its performance, and you should not place undue reliance on such statements. Forward-looking statements are subject to many risks, uncertainties and other variable circumstances. Many of these risks are outside of the Company's control and could cause its actual results to differ materially from those it thought would occur. The forward-looking statements included in this presentation are made only as of the date hereof. The Company does not undertake, and specifically decline, any obligation to update any such statements or to publicly announce the results of any revisions to any of such statements to reflect future events or developments.

Medivir - helping patients through transformative medicines


- Broad pipeline with four candidate drugs in clinical development
- Focus on cancer indications with a high level of unmet medical need and large commercial potential
- World-wide rights to all programs
- Near-term value inflection points
- Strong management and cost-effective virtual organization

Broad and robust pipeline

PROJECT & MECHANISM	DISEASE AREA	RESEARCH	PRECLINICAL	PHASE I	PHASE II	PHASE III	EXCLUSIVITY
Remetinostat HDAC INHIBITOR (TOPICAL)	Cutaneous T-cell lymphoma (MF)	<div></div>					IP: 2034
	Basal cell carcinoma ¹⁾	<div></div>					
Birinapant SMAC MIMETIC (INTRAVENOUS)	Solid tumors (combo with Keytruda®)	<div></div>					IP: 2034
MIV-818 NUCLEOTIDE DNA POLYMERASE INHIBITOR (ORAL)	Hepatocellular carcinoma	<div></div>					IP: 2035
MIV-828 NUCLEOTIDE DNA POLYMERASE INHIBITOR (INTRAVENOUS)	Hematological malignancies (acute myeloid leukemia)	<div></div>					IP: 2039 est.
MIV-711 CATHEPSIN K INHIBITOR (ORAL)	Osteoarthritis	<div></div>					IP: 2029

¹⁾ Investigator sponsored study at Stanford U.

Remetinostat for early-stage cutaneous T-cell lymphoma




MF-CTCL: orphan blood cancer indication

Cutaneous T-cell lymphoma (CTCL) affects lymphocytes (cells belonging to the immune defense system) located in the skin and typically has a chronic course.

- CTCL is a rare form of non-Hodgkin lymphoma primarily present in the skin. Mycosis fungoides (MF) is the most common form of CTCL
- Annual new cases; US ~ 2,000; EU ~ 3,000; Sweden ~ 25
- Five-year survival: ~ 85%; more than 16,000 US patients live with MF-CTCL
- Skin lesions and severe itching are common and affect patients quality of life
- Early stage disease lasts for long periods and requires well tolerated therapy
- Available treatments, including systemic HDAC inhibitors, have severe side effects

Remetinostat: for treatment of early stage MF-CTCL

- Remetinostat is a histone deacetylase (HDAC) inhibitor
- Remetinostat's unique chemistry and topical formulation provides for activity in skin and rapid degradation in blood
- Approved HDAC inhibitors not used in early-stage MF-CTCL patients
- US orphan drug designation



Remetinostat: clinical Proof-of-Concept phase II MF-CTCL study

Twelve months phase II data shows reduction in both lesions and severe itch

Dose	1% 1x/day n=20	0.5% 2x/day n=20	1% 2x/day n=20
Lesion responses ¹	20%	25%	40%
Patients with clinically significant pruritus	1% 1x/day n=8/20 (40%)	0.5% 2x/day n=6/20 (30%)	1% 2x/day n=10/20 (50%)
Pruritus responses	37.5%	50%	80%

Well tolerated:


- No HDAC inhibitor-associated systemic adverse events
- Median time on treatment: 332 days (1% 2x/day dose)

1) Confirmed responses based on CAILS, the Composite Assessment of Index Lesion Severity

Remetinostat: next steps

- End of phase II discussions with the FDA to define phase III design
- Approvals in MF-CTCL have been based on pivotal clinical studies involving less than 400 patients
- One phase III study expected to be sufficient for NDA
- Phase III study will enroll treatment-experienced patients

Birinapant: Uniquely potent against selected solid tumors



Solid tumors: large unmet medical needs

Many patients with solid tumors have few or no options and are in need of effective medicines to extend life. The immuno-oncology medicine Keytruda® on its own is not sufficiently effective in treatment of certain solid tumors.

Colorectal cancer indication (CRC)

- The second most common cancer in women and the third in men
- Estimated new cases 2018: US: ~ 140,000; EU: ~ 490,000; Sweden: ~ 6,200
- Five-year survival: 14%

Other cancer indications


- Ovarian cancer, the leading cause of mortality due a gynecologic tumor
 - Estimated new cases 2018: US: ~ 22,000; EU: ~ 23,000 Sweden: ~ 700
 - Five-year survival: 47%
- Cervical cancer, the third most common cancer in women world-wide
 - Estimated new cases 2018: US: ~ 13,000; EU: ~ 60,000; Sweden: ~ 450
 - Five-year survival: 62.5%

Birinapant may benefit patients with inadequate response to immuno-oncology therapies

- Birinapant, a SMAC mimetic, enables tumor cell death and augments the immune system
- Great potential to improve treatment of cancers when combined with immuno-therapy
- Ongoing collaboration with Merck for a phase I/II study in solid tumors
 - Joint development committee oversees the study
 - Keytruda® provided at no cost by Merck
 - Medivir retains full global rights to birinapant and data


Birinapant/Keytruda[®] combination - phase I/II study ongoing

- Dose escalation fully enrolled; Interim analysis October 8, 2018: n=12
 - One CRC patient has achieved partial response, which had been maintained for 44 weeks
 - Two patients had stable disease for 18 weeks
 - Safety and tolerability: No concerns



- Next steps: evaluation of dose escalation study and determination of phase II dose

MIV-818: Nucleotide prodrug for the treatment of liver cancer



Liver cancer focus: hepatocellular carcinoma (HCC) and intrahepatic cholangiocarcinoma

- HCC is the third leading cause of cancer-related deaths worldwide
 - Estimated new cases 2018: Asia: ~ 610,000; US: ~ 42,000; EU: ~ 82,000; Sweden: ~ 550
 - Orphan disease in Western markets, but one of fastest growing and most deadly cancers in US
 - High incidence in Asia including China - Hepatitis B & C very common
 - Five-year survival: 18%
 - Genetically heterogeneous leading to limited effect of molecularly targeted therapies
- Intrahepatic cholangiocarcinoma is the second most common primary liver tumor
 - Medium survival is only twelve months
- Existing treatment options provide very little survival benefit

MIV-818: prodrug for enhanced efficacy and safety in liver cancer (HCC) therapy


Troxacitabine

- Clinically active but failed due to systemic dose-limiting toxicities




MIV-818

- Enhanced activity
- Selectivity for cancer
- Improved delivery to the liver
- Oral administration
- Limited systemic side effect



MIV-828: For acute myeloid leukemia



MIV-828: Summary

Acute myeloid leukemia (AML) leads to an overproduction of abnormal immature cells in the bone marrow and prevents development of normal blood cells, which could lead to anemia, bleeding episodes and a high susceptibility for severe infections.

- Expected new cases 2018: US: ~ 19,500; EU: ~ 43,000; Sweden: ~ 350
- Affects all ages, median age at diagnosis 68 years in the US
- Five-year survival: 50% in younger patients (<60y) and less than 10% in patients > 70y


Opportunity in hematological cancers

- Better tolerated and more effective agent in patients with AML and other hematological cancers
- Initial development in relapsed/refractory AML patients

Profile of MIV-828

- Nucleotide prodrug based on one of Medivir's proprietary areas of expertise
- Active metabolite shown to have potent anti-leukemic activity in preclinical *in vivo* models
- Increased activity and reduced susceptibility to pharmacologic resistance mechanisms of prodrugs
- Preclinical activity also demonstrated against T-cell lymphoma


MIV-711: Cathepsin K inhibitor with FDA fast track status



Osteoarthritis (OA): the most common form of joint disease

- Affects >30m adults in the US, and ~240m worldwide
- Prevalence increasing due to aging population and obesity epidemic
- Current treatments focus only on pain relief
- Large unmet medical need for a disease-modifying drug (DMOAD) with potential to slow, halt or reverse the progression of OA

Cathepsin K protease is involved in the breakdown of collagen I in both bone and collagen II in cartilage



MIV-711: positive effects on joint structure and signals of benefit on clinical symptoms

- Study design (MIV-711-201): 3 arms, 26 weeks treatment
- Study design (MIV-711-202): 200 mg MIV-711, 26 additional weeks

MIV-711-201: Change from baseline vs week 26


	PBO n=80	n=80 100 mg MIV-711 QD	n=80 200 mg MIV-711 QD
Femur bone area (mm ²)	23.2	8.1	8.2
Cartilage thickness (mm)	-0.066	0.008	-0.017

- A trend consistently favoring MIV-711 arms in all predefined analyses of clinical outcomes, e.g. knee pain and knee function
- Safety and tolerability profile supporting advancement of MIV-711 as a disease-modifying OA drug candidate

New US FDA guidelines in OA may enable pathway for accelerated regulatory approval

- 1999 Draft FDA Guidance in OA was withdrawn August 2018
 - Relied on ordinary x-ray as the best accepted marker for effects on structure, leading to unfeasibly large and long clinical trials
- New draft FDA Guidance published August 2018, focused on structural endpoints in OA development
 - Confirms the FDA's view that OA can be a serious disease and that treatments affecting joint structure are missing
 - The FDA recognizes the need for new approaches in the development of DMOADs and welcomes efforts to establish confidence that measures of structural progression can predict functional outcomes
 - Aim is to be able to accept structural endpoints as valid outcome measures for accelerated approval





Corporate information



About Medivir

- Founded in 1988
- Two medicines developed to the market
- Listed on the Nasdaq Stockholm Main Board
- Market cap as of Nov 15, 2018: approximately SEK 750 million
- Cash position as of Sept 30, 2018: SEK 357.1 million
- Located in Huddinge, Sweden

Strong Management Team



Near term value inflection points

- Birinapant/Keytruda® : completion phase I study – Q4 2018
- Remetinostat: FDA advice on phase III study design – Q1 2019
- MIV-818: completion phase Ia study – Q2 2019